

Name: \_\_\_\_\_  
 Date of birth: \_\_\_\_\_

For use with **EpiPen®** adrenaline (epinephrine) autoinjectors



Confirmed allergens:  
 \_\_\_\_\_

Family/emergency contact name(s):  
 1. \_\_\_\_\_

Mobile Ph: \_\_\_\_\_

2. \_\_\_\_\_

Mobile Ph: \_\_\_\_\_

Plan prepared by doctor or nurse practitioner (np):  
 \_\_\_\_\_

The treating doctor or np hereby authorises medications specified on this plan to be given according to the plan, as consented by the patient or parent/guardian.

Whilst this plan does not expire, review is recommended by DD/MM/YY

Signed: \_\_\_\_\_

Date: \_\_\_\_\_

## SIGNS OF MILD TO MODERATE ALLERGIC REACTION

- Swelling of lips, face, eyes
- Tingling mouth
- Hives or welts
- Abdominal pain, vomiting - **these are signs of anaphylaxis for insect allergy**

## ACTION FOR MILD TO MODERATE ALLERGIC REACTION

- For insect allergy - flick out sting if visible
- For tick allergy  seek medical help or  freeze tick and let it drop off
- Stay with person, call for help and locate adrenaline autoinjector
- Give antihistamine (if prescribed) \_\_\_\_\_
- Phone family/emergency contact

**Mild to moderate allergic reactions (such as hives or swelling) may not always occur before anaphylaxis**

## WATCH FOR ANY ONE OF THE FOLLOWING SIGNS OF ANAPHYLAXIS (SEVERE ALLERGIC REACTION)

- Difficult or noisy breathing
- Difficulty talking or hoarse voice
- Swelling of tongue
- Persistent dizziness or collapse
- Swelling or tightness in throat
- Pale and floppy (young children)
- Wheeze or persistent cough

## ACTION FOR ANAPHYLAXIS

- 1 LAY PERSON FLAT - do NOT allow them to stand or walk**
- If unconscious or pregnant, place in recovery position - on left side if pregnant, as shown below**
- If breathing is difficult allow them to sit with legs outstretched**
- Hold young children flat, not upright**



- 2 GIVE ADRENALINE AUTOINJECTOR**
- 3 Phone ambulance - 000 (AU) or 111 (NZ)**
- 4 Phone family/emergency contact**
- 5 Further adrenaline may be given if no response after 5 minutes**
- 6 Transfer person to hospital for at least 4 hours of observation**

**IF IN DOUBT GIVE ADRENALINE AUTOINJECTOR**  
**Commence CPR at any time if person is unresponsive and not breathing normally**

**ALWAYS GIVE ADRENALINE AUTOINJECTOR FIRST, and then asthma reliever puffer** if someone with known asthma and allergy to food, insects or medication has **SUDDEN BREATHING DIFFICULTY** (including wheeze, persistent cough or hoarse voice) even if there are no skin symptoms

Asthma reliever medication prescribed:  Y  N

*Note: If adrenaline is accidentally injected (e.g. into a thumb) phone your local poisons information centre. Continue to follow this action plan for the person with the allergic reaction.*


### How to give EpiPen®

- Form fist around EpiPen® and **PULL OFF BLUE SAFETY RELEASE**
- Hold leg still and **PLACE ORANGE END** against outer mid-thigh (with or without clothing)
- PUSH DOWN HARD** until a click is heard or felt and hold in place for 3 seconds **REMOVE EpiPen®**

EpiPen® is prescribed as follows:  
 • EpiPen® Jr (150 mcg) for children 7.5-20kg  
 • EpiPen® (300 mcg) for children over 20kg and adults

# ACTION PLAN FOR Anaphylaxis

Name: \_\_\_\_\_ For use with **Anapen®** adrenaline (epinephrine) autoinjectors  
 Date of birth: \_\_\_\_\_



Confirmed allergens:

\_\_\_\_\_

Family/emergency contact name(s):

1. \_\_\_\_\_

Mobile Ph: \_\_\_\_\_

2. \_\_\_\_\_

Mobile Ph: \_\_\_\_\_

Plan prepared by doctor or nurse practitioner (np): \_\_\_\_\_

The treating doctor or np hereby authorises medications specified on this plan to be given according to the plan, as consented by the patient or parent/guardian.

Whilst this plan does not expire, review is recommended by \_\_\_\_\_

Signed: \_\_\_\_\_

Date: \_\_\_\_\_

## SIGNS OF MILD TO MODERATE ALLERGIC REACTION

- Swelling of lips, face, eyes
- Hives or welts
- Tingling mouth
- Abdominal pain, vomiting - **these are signs of anaphylaxis for insect allergy**

## ACTION FOR MILD TO MODERATE ALLERGIC REACTION

- For insect allergy - flick out sting if visible
- For tick allergy  seek medical help or  freeze tick and let it drop off
- Stay with person, call for help and locate adrenaline autoinjector
- Give antihistamine (if prescribed) \_\_\_\_\_
- Phone family/emergency contact

**Mild to moderate allergic reactions (such as hives or swelling) may not always occur before anaphylaxis**

## WATCH FOR ANY ONE OF THE FOLLOWING SIGNS OF ANAPHYLAXIS (SEVERE ALLERGIC REACTION)

- Difficult or noisy breathing
- Swelling of tongue
- Swelling or tightness in throat
- Wheeze or persistent cough
- Difficulty talking or hoarse voice
- Persistent dizziness or collapse
- Pale and floppy (young children)

## ACTION FOR ANAPHYLAXIS

- LAY PERSON FLAT - do NOT allow them to stand or walk**
  - If unconscious or pregnant, place in recovery position - on left side if pregnant, as shown below
  - If breathing is difficult allow them to sit with legs outstretched
  - Hold young children flat, not upright



- GIVE ADRENALINE AUTOINJECTOR**
- Phone ambulance - 000 (AU) or 111 (NZ)**
- Phone family/emergency contact**
- Further adrenaline may be given if no response after 5 minutes**
- Transfer person to hospital for at least 4 hours of observation**

### IF IN DOUBT GIVE ADRENALINE AUTOINJECTOR

**Commence CPR at any time if person is unresponsive and not breathing normally**

**ALWAYS GIVE ADRENALINE AUTOINJECTOR FIRST, and then asthma reliever puffer** if someone with known asthma and allergy to food, insects or medication has **SUDDEN BREATHING DIFFICULTY** (including wheeze, persistent cough or hoarse voice) even if there are no skin symptoms

Asthma reliever medication prescribed:  Y  N

*Note: If adrenaline is accidentally injected (e.g. into a thumb) phone your local poisons information centre. Continue to follow this action plan for the person with the allergic reaction.*

### How to give Anapen®



**1**  
PULL OFF BLACK NEEDLE SHIELD



**2**  
PULL OFF GREY SAFETY CAP from red button



**3**  
PLACE NEEDLE END FIRMLY against outer mid-thigh at 90° angle (with or without clothing)



**4**  
PRESS RED BUTTON so it clicks and hold for 10 seconds. REMOVE Anapen®

Anapen® is prescribed as follows:

- Anapen® 150 Junior for children 7.5-20kg
- Anapen® 300 for children over 20kg and adults
- Anapen® 500 for children and adults over 50kg